



PREGNANT WOMEN SMOKING RATES BY INDIANA COUNTIES

Percent of mothers who reported smoking during pregnancy

County	Percent	County	Percent	County	Percent	County	Percent
Adams	11.6	Franklin	25.5	Lawrence	25.6	Rush	25.1
Allen	18.1	Fulton	28.6	Madison	26.7	St. Joseph	15.9
Bartholomew	13.5	Gibson	24.4	Marion	19.7	Scott	38.1
Benton	21.3	Grant	31.8	Marshall	21.2	Shelby	28.4
Blackford	30.3	Greene	23.3	Martin	18.8	Spencer	26.4
Boone	14.4	Hamilton	6.8	Miami	27.9	Starke	31.8
Brown	20.5	Hancock	18.5	Monroe	20.2	Steuben	26.2
Carroll	18.2	Harrison	22.0	Montgomery	28.4	Sullivan	26.5
Cass	23.8	Hendricks	13.4	Morgan	25.5	Switzerland	31.4
Clark	24.6	Henry	26.0	Newton	25.2	Tippecanoe	13.8
Clay	28.2	Howard	23.4	Noble	26.6	Tipton	19.2
Clinton	19.9	Huntington	23.8	Ohio	27.9	Union	13.5
Crawford	34.9	Jackson	23.6	Orange	25.5	Vanderburgh	23.7
Davies	20.7	Jasper	21.3	Owen	29.3	Vermillion	31.8
Dearborn	33.7	Jay	20.6	Parke	27.9	Vigo	26.0
Decatur	34.3	Jefferson	33.8	Perry	38.7	Wabash	24.4
DeKalb	32.7	Jennings	22.1	Pike	20.4	Warren	16.0
Delaware	22.5	Johnson	20.0	Porter	17.5	Warrick	17.6
Dubois	13.4	Knox	32.2	Posey	19.2	Washington	29.0
Elkhart	16.8	Kosciusko	19.2	Pulaski	30.1	Wayne	25.5
Fayette	25.9	LaGrange	12.4	Putnam	24.3	Wells	20.7
Floyd	25.3	Lake	16.2	Randolph	27.7	White	20.5
Fountain	26.0	LaPorte	25.5	Ripley	29.0	Whitley	22.5

- Overall **20.3%** of women in Indiana reported smoking during pregnancy.
- **70%** of Indiana's counties have a smoking during pregnancy rate higher than the Indiana average.
- **12.3%** of women in the United States smoked during pregnancy. (1999)
- All but two Indiana counties have a smoking during pregnancy rates higher than the United States average.

Sources: 2000 Indiana Birth Certificate Data; National Vital Statistics Report, "Smoking During Pregnancy in the 1990s", 2001.

Indiana Tobacco Prevention and Cessation
 150 West Market – Suite 406 – Indianapolis, IN 46204
 phone – 317.234.1787 fax – 317.234.1786
www.itpc.in.gov

updated 5/2002